

THE CLASSIFICATION OF 2-COMPACT GROUPS

JESPER GRODAL

(JOINT WORK WITH KASPER K. S. ANDERSEN)

ABSTRACT. In this talk I'll announce and explain a proof of the classification of 2-compact groups, joint with K. Andersen, hence completing the classification of p -compact groups at all primes p . A p -compact group, as introduced by Dwyer-Wilkerson, is a homotopy theoretic version of a compact Lie group, but with all its structure concentrated at a single prime p . Our classification states that there is a 1-1-correspondence between connected 2-compact groups and root data over the 2-adic integers (which will be defined in the talk). As a consequence we get the conjecture that every connected 2-compact group is isomorphic to a product of the 2-completion of a compact Lie group and copies of the exotic 2-compact group $DI(4)$, constructed by Dwyer-Wilkerson. The major new input in the proof over the proof at odd primes (due to Andersen-Grodal-Møller-Viruel) is a thorough analysis of the concept of a root datum for 2-compact groups and its relationship with the maximal torus normalizer. With these tools in place we are able to produce a proof which to a large extent avoids case-by-case considerations.

This is a summary of my lecture given at the “Conference on Pure and Applied Topology”, Isle of Skye, Friday June 24, 2005. I announced and sketched a proof of the classification of 2-compact groups, in joint work with Kasper Andersen. This will appear in the papers [2] and [3].

Recall that a p -compact group is a triple $(X, BX, e : X \xrightarrow{\cong} \Omega BX)$ where BX is a pointed, connected, p -complete space of the homotopy type of a CW -complex, X satisfies that $H^*(X; \mathbb{F}_p)$ is finite over $\mathbb{F}_p$, and e is a homotopy equivalence. They are homotopy theoretic analogs of compact Lie groups, and were introduced by Dwyer-Wilkerson in [6]. A p -compact group is said to be connected if X is a connected space.

Our main theorem is the following

Theorem 1.1 ([3]). *Let $(X, BX, e : X \xrightarrow{\cong} \Omega BX)$ be a connected 2-compact group. Then*

$$BX \simeq BG_2^{\wedge} \times BDI(4)^s$$

where $BG_2^{\wedge}$ is the 2-completion of a connected compact Lie group G , and $BDI(4)$ is the classifying space of the exotic 2-compact group $DI(4)$ constructed by Dwyer-Wilkerson in [5], $s \geq 0$.

A corresponding statement for odd primes was proved by the authors together with Møller and Viruel in [4]. Partial results for $p = 2$ have been obtained by Dwyer-Miller-Wilkerson, Notbohm, Viruel, Møller, and Vavpetič-Viruel. Our proof is a self-contained induction.

There is a better, more precise, formulation of our theorem which both makes it clear why it is the correct 2-local version of the classification of compact Lie groups and suggests a possible strategy of proof. It is based on the notion of a root datum over the p -adic integers $\mathbb{Z}_p$, which we now introduce.

Partially supported by NSF grant DMS-0354633.

For R a principal ideal domain, an R -root datum $\mathbf{D}$ is defined to be a triple $(W, L, \{Rb_\sigma\})$, where L is a free R -module of finite rank, $W \subseteq \text{Aut}_R(L)$ is a finite subgroup generated by reflections (i.e., elements σ such that $1 - \sigma \in \text{End}_R(L)$ has rank one), and $\{Rb_\sigma\}$ is a collection of rank one submodules of L , indexed by the reflections σ in W , satisfying

$$\text{im}(1 - \sigma) \subseteq Rb_\sigma \subseteq \ker\left(\sum_{i=0}^{|\sigma|-1} \sigma^i\right) \text{ and } w(Rb_\sigma) = Rb_{w\sigma w^{-1}} \text{ for all } w \in W$$

The element $b_\sigma \in L$, called the *coroot* corresponding to σ , is determined up to a unit in R . Together with σ it determines a root $\beta_\sigma : L \rightarrow R$ via the formula

$$\sigma(x) = x + \beta_\sigma(x)b_\sigma$$

If $R = \mathbb{Z}$ then there is a 1-1-correspondence between $\mathbb{Z}$ -root data and classically defined root data, by to $(W, L, \{\mathbb{Z}b_\sigma\})$ associating $(L, L^*, \{\pm b_\sigma\}, \{\pm \beta_\sigma\})$; see [8, Prop. 2.16]. If $R = \mathbb{Z}$ or $\mathbb{Z}_p$, instead of the collection $\{Rb_\sigma\}$ one can equivalently consider their span, the *coroot lattice*, $L_0 = +_\sigma Rb_\sigma \subseteq L$, and this was the definition given in [4, § 1], under the name “ R -reflection datum”. Two R -root data $\mathbf{D} = (W, L, \{Rb_\sigma\})$ and $\mathbf{D}' = (W', L', \{Rb'_\sigma\})$ are said to be isomorphic if there exists an isomorphism $\varphi : L \rightarrow L'$ such that $\varphi W \varphi^{-1} = W'$ and $\varphi(Rb_\sigma) = Rb'_{\varphi\sigma\varphi^{-1}}$. In particular the automorphism group is given by $\text{Aut}(\mathbf{D}) = \{\varphi \in N_{\text{Aut}_R(L)}(W) \mid \varphi(Rb_\sigma) = Rb_{\varphi(\sigma)}\}$ and we define the outer automorphism group as $\text{Out}(\mathbf{D}) = \text{Aut}(\mathbf{D})/W$.

We now explain how to associate a $\mathbb{Z}_p$ -root datum to a p -compact group. By a theorem of Dwyer-Wilkerson any p -compact group (X, BX, e) has a maximal torus, which is a map $i : BT = (BS^1_p)^r \rightarrow BX$ satisfying that the fiber has finite $\mathbb{F}_p$ -cohomology and non-trivial Euler characteristic. Replacing i by an equivalent fibration, we define the *Weyl space* $\mathcal{W}_X(T)$ as the topological monoid of self-maps $BT \rightarrow BT$ over i . The *Weyl group* is defined as $W_X(T) = \pi_0(\mathcal{W}_X(T))$ and the classifying space of the maximal torus normalizer is defined as $B\mathcal{N}_X(T) = BT_{h\mathcal{W}_X(T)}$. Now, by definition, W_X acts on $L = \pi_2(BT)$ and it is a theorem of Dwyer-Wilkerson that, if X is connected, this gives a faithful representation of W_X in $\text{Aut}_{\mathbb{Z}_p}(L)$ as a finite $\mathbb{Z}_p$ -reflection group. There is also an easy formula for the b_σ in terms of the maximal torus normalizer $\mathcal{N}_X$, for which we refer to [8], [2], or [3]. This explains how to associate a $\mathbb{Z}_p$ -root datum $\mathbf{D}_X$ to a connected p -compact group X . (In the case where p is odd the root datum is in fact completely determined by the finite $\mathbb{Z}_p$ -reflection group (W, L) , which explains the formulation of our classification theorem with Møller and Viruel in [4].)

We are now ready to state the precise version of our main theorem.

Theorem 1.2 ([3]). *The assignment which to a connected 2-compact group X associates its $\mathbb{Z}_2$ -root datum $\mathbf{D}_X$ root gives a one-to-one correspondence between connected 2-compact groups and $\mathbb{Z}_2$ -root data. Furthermore the map $\Phi : \pi_0(\text{Aut}(BX)) \rightarrow \text{Out}(\mathbf{D}_X)$ is an isomorphism, and $B\text{Aut}(BX)$ is the unique total space of a split fibration*

$$B^2\mathcal{Z}(\mathbf{D}) \rightarrow B\text{Aut}(BX) \rightarrow B\text{Out}(\mathbf{D}_X)$$

Here $\mathcal{Z}(\mathbf{D})$ is the center of the root datum $\mathbf{D}$, defined so as to agree with the formula for the center of a p -compact group given in [7], and $B\text{Aut}(BX)$ denotes the classifying space of the topological monoid of self-homotopy equivalences of BX .

We remark that since we have completely described connected p -compact groups *as well as their space of automorphisms*, we also get a classification for non-connected 2-compact groups. (The situation here is totally analogous to the case of compact Lie groups.) If X

is a connected p -compact group with a given root datum $\mathbf{D}$ and π is a given p -group, then the p -compact groups with connected component isomorphic to X and component group π are in 1-1-correspondence with the $\text{Out}(\pi)$ -orbits of the set $[B\pi, B\text{Aut}(BX)]$. Since both X and $B\text{Aut}(BX)$ is completely described in terms of $\mathbf{D}_X$ by the main theorem, this gives a classification also in the non-connected case. We refer to [3] for a more detailed description.

The main theorem has a number of corollaries. The most important is perhaps that it gives a proof of the maximal torus conjecture, giving a purely homotopy theoretic characterization of compact Lie groups amongst finite loop spaces.

Theorem 1.3 (Maximal torus conjecture [3]). *The classifying space functor, which to a compact Lie group G associates the finite loop space $(G, BG, e : G \xrightarrow{\cong} \Omega BG)$ gives a one-to-one correspondence between compact Lie groups and finite loop spaces with a maximal torus. (Moreover, if G is connected we have a split fibration $B^2\mathcal{Z}(\mathbf{D}_G) \rightarrow B\text{Aut}(BG) \rightarrow B\text{Out}(\mathbf{D}_G)$.)*

The fact that the functor “ B ” is faithful was already known by work of Notbohm, Møller, and Osse and the statement about the space $B\text{Aut}(BG)$ follows easily from earlier work of Jackowski-McClure-Oliver and Dwyer-Wilkerson. The new, and a priori quite surprising, result here is the statement that if a finite loop space has a maximal torus, then it has to come from a compact Lie group.

Another application of the classification of 2-compact groups is to give an answer to the so-called Steenrod problem for $p = 2$ (see [13] and [12]), which asks which graded polynomial algebras can occur as the mod 2 cohomology ring of a space? Steenrod’s problem was solved for p “large enough” by Adams-Wilkerson [1] and for all odd primes by Notbohm [11] using a partial classification of p -compact groups, p odd.

Theorem 1.4 (Steenrod’s problem for $p = 2$ [3]). *Suppose that P^* is a graded polynomial algebra over $\mathbb{F}_2$ in finitely many variables. If P^* occurs as $H^*(Y; \mathbb{F}_2)$ for some space Y (no assumptions), then P^* is isomorphic, as a graded algebra, to*

$$H^*(BG; \mathbb{F}_2) \otimes H^*(BDI(4); \mathbb{F}_2)^{\otimes s} \otimes Q^*$$

where G is a connected semi-simple Lie group and Q^* is a polynomial ring with generators in degrees one and two.

In particular if P^* is assumed to have generators in degree ≥ 3 then G has to be simply connected and P^* is a tensor product of the following graded algebras:

$$\begin{array}{ll} \mathbb{F}_2[x_4, x_6, \dots, x_{2n}] & (\text{SU}(n)) \\ \mathbb{F}_2[x_4, x_8, \dots, x_{4n}] & (\text{Sp}(n)) \\ \mathbb{F}_2[x_4, x_6, x_7, x_8] & (\text{Spin}(7)) \\ \mathbb{F}_2[x_4, x_6, x_7, x_8, x'_8] & (\text{Spin}(8)) \\ \mathbb{F}_2[x_4, x_6, x_7, x_8, x_{16}] & (\text{Spin}(9)) \\ \mathbb{F}_2[x_4, x_6, x_7] & (G_2) \\ \mathbb{F}_2[x_4, x_6, x_7, x_{16}, x_{24}] & (F_4) \\ \mathbb{F}_2[x_8, x_{12}, x_{14}, x_{15}] & (DI(4)) \end{array}$$

It seems reasonable that one can in fact list all polynomial rings which occur as $H^*(BG; \mathbb{F}_2)$ for G semi-simple, although we have not been able to locate such a list in the literature; for G simple a list can be found in [10].

We finally point out that many classical theorems from Lie theory by Borel, Bott, Demazure, and others also carry over to 2-compact groups via the classification. Applications of this type were already pointed out in [4], to which we refer.

We end this short summary by saying a few words about the proofs. Our strategy is an elaboration of the strategy we pursued with Møller and Viruel in [4], but with several new additions, most importantly a development and utilization of the theory of p -adic root data. The sketch below is intended to sum up the main steps without introducing too much notation. We focus on the uniqueness statement, that two connected p -compact groups with isomorphic root data are isomorphic.

It is a recent theorem of Dwyer-Wilkerson [8], extending old work of Tits, that the maximal torus normalizer $\mathcal{N}_X$ in a 2-compact group X can be recovered from the root datum $\mathbf{D}_X$. However, $\mathcal{N}_X$ has a larger group of automorphisms than X , which means that it is not the right classifying invariant.

This problem can be overcome, by also keeping track of certain “root subgroups”, a machinery which we set up in [2]. In a given maximal torus normalizer $\mathcal{N}$, for each reflection σ and coroot b_σ one can algebraically construct a root subgroup $\mathcal{N}_\sigma$ of $\mathcal{N}$. In the setting of algebraic groups, this “root subgroup” $\mathcal{N}_\sigma$ will be the maximal torus normalizer of $\langle U_\alpha, U_{-\alpha} \rangle$, where U_α is the root subgroup in the sense of algebraic groups corresponding to the root α dual to the coroot b_σ .

Theorem 1.5 ([2]). *Let X be a connected 2-compact group. The canonical map $\text{Out}(BX) \rightarrow \text{Out}(\mathbf{D}_X)$ factors*

$$\Phi : \text{Out}(BX) \rightarrow \text{Out}(B\mathcal{N}, \{B\mathcal{N}_\sigma\}) \xrightarrow{\cong} \text{Out}(\mathbf{D}_X)$$

and we likewise have a canonical map

$$\Phi : B\text{Aut}(BX) \rightarrow B\text{aut}(\mathbf{D}_X)$$

where $B\text{aut}(\mathbf{D}_X)$ is a certain space defined in [2] refining $B\text{Aut}(B\mathcal{N}, \{B\mathcal{N}_\sigma\})$.

These maps will be shown to be isomorphisms inductively, as part of proving Theorem 1.2. Since the pair $(\mathcal{N}, \{\mathcal{N}_\sigma\})$ carries the same structure and automorphisms as $\mathbf{D}$, but is closer to X , it is more useful than $\mathbf{D}$ for classification purposes.

Assume that we have two connected p -compact groups X and X' with the same root datum $\mathbf{D}$ and hence the same maximal torus normalizer and root subgroups $(\mathcal{N}, \{\mathcal{N}_\sigma\})$, and assume by induction that Theorem 1.2 is true for all connected p -compact groups of smaller cohomological dimension. The first step is to observe that one can reduce to the case where X and X' are simple and center-free, mimicking the proofs in [4] but everywhere keeping track of the root subgroups. So we can assume that we are in the following situation

$$\begin{array}{ccc} & (B\mathcal{N}, \{B\mathcal{N}_\sigma\}_\sigma) & \\ \swarrow & & \searrow \\ BX & \text{-----} & BX' \end{array}$$

where the dotted arrow is the one that we want to construct.

Using that every element of order p can be conjugated into the maximal torus, uniquely up to conjugation in $\mathcal{N}$, we can by induction construct the dotted arrow on the connected

component of the centralizer of every element $\nu : B\mathbb{Z}/p \rightarrow BX$ of order p in X .

$$\begin{array}{ccc} & (BC_{\mathcal{N}}(\nu)_1, \{BC_{\mathcal{N}}(\nu)_{1\sigma}\}_{\sigma}) & \\ & \swarrow \quad \searrow & \\ BC_X(\nu)_1 & \xrightarrow{\quad\quad\quad} & BC_{X'}(\nu)_1 \end{array}$$

Using that we have control of the whole *space* of self-equivalences of $BC_X(\nu)_1$ by Theorem 1.5 and induction, one sees that this map is equivariant with respect to the component group and hence extends uniquely to the whole centralizer $BC_X(\nu)$.

Now for a general elementary abelian subgroup $\nu : E \rightarrow X$ of X we can pick an element of order p in E , and we get via restriction a map

$$BC_X(\nu) \rightarrow BC_X(\mathbb{Z}/p) \rightarrow BC_{X'}(\mathbb{Z}/p) \rightarrow BX'$$

To make sure that these maps are chosen in a compatible way, one has to show that this map does not depend on the choice of rank one subgroup of E . We developed some techniques in [4] for handling this kind of situation. Using easy (one page) case-by-case arguments on the level of 2-adic root data, we are able to verify that the assumptions of [4] are always satisfied.

Since these maps from centralizers are chosen in a compatible way, they combine to form an element

$$\vartheta \in \lim_{\nu \in \mathbf{A}(X)}^0 [BC_X(\nu), BX']$$

where $\mathbf{A}(X)$ is the Quillen category of X . Furthermore the construction of this element allows one to show that X and X' have the same p -fusion and in particular the same maximal torus p -normalizer and the same $\mathbb{Z}_p$ -cohomology. We are left with a rigidification question.

One approach could be to show that the relevant obstruction groups

$$\lim_{\vartheta \in \mathbf{A}(X)}^* \pi_*(\text{map}(BC_X(\nu), BX')_{[\vartheta]})$$

vanish, which would show that our diagram rigidifies and produces an equivalence

$$BX \simeq \text{hocolim}_{\nu \in \mathbf{A}(X)} BC_X(\nu) \xrightarrow{\sim} BX'$$

It is immediate to see that this is the case for many groups X including $DI(4)$, and it was the approach taken for p odd in [4]. There we for instance also show that it holds for $X = \text{PU}(n)\hat{2}$, by the same calculation as for p odd, but it is more difficult to verify e.g., for the exceptional groups.

We however take a different approach: Since $DI(4)$ is easily dealt with, we can assume that $BX = BG\hat{2}$ for some simple, center-free Lie group. For each p -radical subgroup P of G we have a map $BP \rightarrow BC_G({}_pZ(P))_{\hat{p}} \rightarrow BX'$. Since X and X' have the same $\mathbb{Z}_p$ -homology, and in particular the same fundamental group, we can lift this map to a map $B\tilde{P} \rightarrow B\tilde{X}'$, where $\tilde{P}$ is the preimage of P in $\tilde{G}$, the universal cover of G , and $\tilde{X}'$ is the universal cover of X' .

One now verifies that these maps assemble to give an element in

$$\lim_{\tilde{G}/\tilde{P} \in \mathbf{O}_p^r(\tilde{G})}^0 [B\tilde{P}, B\tilde{X}']$$

where $\mathbf{O}_p^r(\tilde{G})$ is the full subcategory of the p -orbit category of $\tilde{G}$ with objects $\tilde{G}/\tilde{P}$ for $\tilde{P}$ a p -radical subgroup of $\tilde{G}$. The obstructions to rigidifying this to get a map

$$B\tilde{G} \simeq \operatorname{hocolim}_{\tilde{G}/\tilde{P} \in \mathbf{O}_p^r(\tilde{G})} B\tilde{P} \rightarrow B\tilde{X}'$$

can be shown to lie in groups which identify with

$$\lim_{\tilde{G}/\tilde{P} \in \mathbf{O}_p^r(\tilde{G})} {}^*\pi_*(Z(\tilde{P}))$$

These groups have been shown to identically vanish by earlier work of Jackowski-McClure-Oliver [9]. Hence the map exists and it now follows easily by the construction that it is an equivalence. Passing to a quotient we get the wanted equivalence $BG \rightarrow BX'$, finishing the proof of the theorem.

REFERENCES

- [1] J. F. Adams and C. W. Wilkerson. Finite H -spaces and algebras over the Steenrod algebra. *Ann. of Math. (2)*, 111(1):95–143, 1980. (Erratum: *Ann. of Math. (2)* **113** (1981), no. 3, 621–622).
- [2] K. K. S. Andersen and J. Grodal. Automorphisms of root data, maximal torus normalizers, and p -compact groups. Preprint 2005. 19 pages.
- [3] K. K. S. Andersen and J. Grodal. The classification of 2-compact groups. in preparation.
- [4] K. K. S. Andersen, J. Grodal, J. M. Møller, and A. Viruel. The classification of p -compact groups for p odd. *Ann. of Math.* (to appear). arXiv:math.AT/0302346. 92 pages.
- [5] W. G. Dwyer and C. W. Wilkerson. A new finite loop space at the prime two. *J. Amer. Math. Soc.*, 6(1):37–64, 1993.
- [6] W. G. Dwyer and C. W. Wilkerson. Homotopy fixed-point methods for Lie groups and finite loop spaces. *Ann. of Math. (2)*, 139(2):395–442, 1994.
- [7] W. G. Dwyer and C. W. Wilkerson. The center of a p -compact group. In *The Čech centennial (Boston, MA, 1993)*, pages 119–157. Amer. Math. Soc., Providence, RI, 1995.
- [8] W. G. Dwyer and C. W. Wilkerson. Normalizers of tori. *Geom. Topol.*, 9:1337–1380 (electronic), 2005.
- [9] S. Jackowski, J. McClure, and B. Oliver. Homotopy classification of self-maps of BG via G -actions. I+II. *Ann. of Math. (2)*, 135(2):183–226, 227–270, 1992.
- [10] A. Kono. On cohomology mod 2 of the classifying spaces of non-simply connected classical Lie groups. *J. Math. Soc. Japan*, 27:281–288, 1975.
- [11] D. Notbohm. Spaces with polynomial mod- p cohomology. *Math. Proc. Cambridge Philos. Soc.*, 126(2):277–292, 1999.
- [12] N. Steenrod. Polynomial algebras over the algebra of cohomology operations. In *H-spaces (Actes Réunion Neuchâtel, 1970)*, Lecture Notes in Mathematics, Vol. 196, pages 85–99. Springer, Berlin, 1971.
- [13] N. E. Steenrod. The cohomology algebra of a space. *Enseignement Math. (2)*, 7:153–178 (1962), 1961.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AARHUS, DK-8000 ÅRHUS C, DENMARK
E-mail address: kksa@math.ku.dk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, IL 60637, USA
E-mail address: jg@math.uchicago.edu